

## Math 2374 - Quiz 5

Name: \_\_\_\_\_

Section: \_\_\_\_\_

Consider the vector field

$$V(x, y) = (2xy^3 + 2x, 3x^2y^2 + \cos y)$$

1. Show that the *scalar curl* is 0. Remember that the scalar curl is the  $z$ -component of the curl of the vector field  $W$  given by

$$W(x, y, z) = (2xy^3 + 2x, 3x^2y^2 + \cos y, 0).$$

**Solution:** The scalar curl is  $\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}$ . One easily sees that  $\partial Q/\partial x = 6xy^2 = \partial P/\partial y$ , so

$$\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} = 6xy^2 - 6xy^2 = 0.$$

Rubric: out of a total of 10 points:

- -1 point for non-trivializing arithmetic mistakes.
- Full credit if they correctly use  $W$ .
- Full credit if they just use the formula.

2. Let  $c$  be a positively oriented simple closed curve in the plane. Show that

$$\int_c V \cdot ds = 0.$$

*You are not given the precise form of the curve, so you may want to avoid integrating directly.*

**Solution:** By Green's theorem we know that

$$\int_c V \cdot ds = \iint_R \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} dx dy,$$

where  $R$  is the region enclosed by the  $c$ . By the above problem the integrand is 0, so we're done.

Rubric: out of a total of 40 points:

- Full credit if they say that there must exist a potential (or they use the potential from the next question).

3. In the last part of the quiz we're going to find a potential  $f$  whose gradient is  $V$ . In other words, we're looking for a function  $f$  satisfying

$$V = \nabla f.$$

1. If such an  $f$  were to exist, it would necessarily need to verify

$$\frac{\partial f(x, y)}{\partial x} = 2xy^3 + 2x.$$

Find a function that satisfies this. Remember to add a  $+C(y)$  at the end, this way you will get *all* such functions.

**Solution:** To find a function whose derivative is a given function we integrate it:

$$f(x, y) = \int 2xy^3 + 2x \, dx = x^2y^3 + x^2 + C(y).$$

2. In the previous question you found *all* functions whose partial derivative with respect to  $x$  is the first component of  $V$ , they have the form

$$f(x, y) = \text{Something} + C(y).$$

You want  $f$  to *also* satisfy

$$\frac{\partial f(x, y)}{\partial y} = 3x^2y^2 + \cos y.$$

Plug the expression of  $f$  that you found above into this equation in order to see what  $C$  needs to look like, and then find a  $C$  which works.

**Solution:** Plugging the expression above we have

$$\frac{\partial}{\partial y}(x^2y^3 + x^2 + C(y)) = 3x^2y^2 + \cos y.$$

The left hand side is

$$3x^2y^2 + C'(y),$$

so the term  $3x^2y^2$  cancels and we get that  $C'(y) = \cos y$ .

We just need to find a function  $C$  that satisfies  $C'(y) = \cos y$ , so  $C(y) = \sin y$ .

Putting it all together we get that a possible  $f$  is

$$f(x, y) = x^2y^3 + x^2 + \sin y.$$

Rubric: out of a total of 50 points:

- Each part is worth 25 points.
- -1 point for non-trivializing arithmetic mistakes.